

# The two-process model of sleep regulation: a reappraisal

ALEXANDER A. BORBÉLY<sup>1</sup>, SERGE DAAN<sup>2</sup>, ANNA WIRZ-JUSTICE<sup>3</sup> and TOM DEBOER<sup>4</sup>

<sup>1</sup>Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland; <sup>2</sup>Centre for Behaviour and Neuroscience, University of Groningen, Groningen, the Netherlands; <sup>3</sup>Centre for Chronobiology, University of Basel Psychiatric Clinics, Basel, Switzerland; <sup>4</sup>Laboratory for Neurophysiology, Department of Molecular Cell Biology, Leiden University Medical Center, Leiden, the Netherlands

## Keywords

rest-activity cycle, napping, circadian phase, forced desynchrony, neuronal light response, seasonal affective disorder

## Correspondence

Alexander A. Borbély, Institute of Pharmacology and Toxicology, University of Zurich, Switzerland.

Tel.: +4144 635 5960;

Fax: +4144 635 5888;

e-mail: borbely@pharma.uzh.ch

Accepted in revised form 5 November 2015;  
received 5 November 2015

DOI: 10.1111/jsr.12371

## SUMMARY

In the last three decades the two-process model of sleep regulation has served as a major conceptual framework in sleep research. It has been applied widely in studies on fatigue and performance and to dissect individual differences in sleep regulation. The model posits that a homeostatic process (Process S) interacts with a process controlled by the circadian pacemaker (Process C), with time-courses derived from physiological and behavioural variables. The model simulates successfully the timing and intensity of sleep in diverse experimental protocols. Electrophysiological recordings from the suprachiasmatic nuclei (SCN) suggest that S and C interact continuously. Oscillators outside the SCN that are linked to energy metabolism are evident in SCN-lesioned arrhythmic animals subjected to restricted feeding or methamphetamine administration, as well as in human subjects during internal desynchronization. In intact animals these peripheral oscillators may dissociate from the central pacemaker rhythm. A sleep/fast and wake/feed phase segregate antagonistic anabolic and catabolic metabolic processes in peripheral tissues. A deficiency of Process S was proposed to account for both depressive sleep disturbances and the antidepressant effect of sleep deprivation. The model supported the development of novel non-pharmacological treatment paradigms in psychiatry, based on manipulating circadian phase, sleep and light exposure. In conclusion, the model remains conceptually useful for promoting the integration of sleep and circadian rhythm research. Sleep appears to have not only a short-term, use-dependent function; it also serves to enforce rest and fasting, thereby supporting the optimization of metabolic processes at the appropriate phase of the 24-h cycle.

## INTRODUCTION

The two-process model of sleep regulation was proposed more than three decades ago. It had a large impact on sleep research and is still a prevalent conceptual model. This is reflected by the persistently high number of citations per year of the original papers. AAB and SD published the original

papers of the two process model (Borbély, 1982; Daan *et al.*, 1984), whereas AAB and AWJ related the model to sleep in depression and the antidepressant effect of sleep deprivation (Borbély and Wirz-Justice, 1982). The present review was triggered by a symposium at the European Sleep Research Society (ESRS) Congress in Tallinn in September 2014, in which the authors highlighted important developments and

assessed merits and limitations of the model.<sup>1</sup> The review focuses on conceptual issues and does not cover other important aspects, such as applications of the model to work schedules or performance measures, its mathematical refinements, age-related changes and new insights into the molecular mechanisms underlying the two processes.

## THE ORIGINAL TWO-PROCESS MODEL OF SLEEP REGULATION AND ITS PREDICTIVE POWER

### A brief restatement of the model


The two-process model posits that the interaction of a homeostatic process depending on sleep and wake (Process S) with a process controlled by the circadian pacemaker (Process C) determines salient aspects of sleep regulation. Process S, representing sleep debt, increases during wakefulness and declines during sleep, within a value-range that oscillates with a periodicity that is normally entrained to day and night by the circadian pacemaker. When S approaches the range's lower boundary it triggers awakening; near the upper boundary it triggers sleep. Non-rapid eye movement (NREM) sleep electroencephalography (EEG) slow wave activity (SWA) represents the principal marker of S during sleep; theta activity in waking is a marker of the rising limb of S (reviewed by Borbély and Achermann, 1999). Core body temperature and melatonin rhythms are markers of C. In animals rendered arrhythmic by lesioning the suprachiasmatic nuclei (SCN), sleep homeostasis is not disrupted (Tobler *et al.*, 1983; Trachsel *et al.*, 1992). This indicates that the two processes are regulated separately. In humans, the forced desynchrony protocol, an imposed sleep-wake periodicity that is outside the range of entrainment of the circadian pacemaker, results in sleep occurring at different circadian phases. This disentangles the sleep-wake cycle from the circadian rhythm, allowing assessment of their separate influence on sleep and performance variables (Dijk and Czeisler, 1995; Wyatt *et al.*, 1999).

<sup>1</sup>Historical note: in the 1980s AAB, SD and AWJ had established a Zürich-Groningen-Basel (ZGB) research triangle that gave rise to a continuous exchange of students and ideas related to sleep regulation, as well as experiments testing the model. TD was a student in Groningen and Zürich, and now focuses his research on the interaction of the two processes. A further key person in the investigation of sleep and depression was the late Rutger van den Hoofdakker in Groningen. Many of the students and postdoctoral researchers who were trained in the ZGB research triangle have continued their research into the two-process model. They include Irene Tobler, Domien Beersma, Peter Achermann, Derk-Jan Dijk, Kurt Kräuchi, Christian Cajochen, Paul Franken, Hanspeter Landolt, Daniel Brunner, Daniel Aeschbach, Esther Werth, Reto Huber and Vladislav Vyazovskiy. Their enthusiasm and creativity were instrumental in the further development and validation of the model.

### Tests of the model

In humans, the model was found initially to be consistent with the phenomenon of internal desynchronization between sleep-wake and circadian body temperature cycles in the absence of time cues, with circadian oscillations in fatigue during prolonged (3-day) sleep deprivation, with sleep fragmentation during continuous bedrest, with SWA in sleep after different durations of sleep deprivation and with sleep duration in shiftworkers (Daan *et al.*, 1984).

Predictions derived from the model were then tested. NREM sleep SWA during brief daytime naps increased with duration of prior wakefulness (Dijk *et al.*, 1987a). After such naps, SWA in normal night sleep was reduced in quantitative agreement with model predictions (Daan *et al.*, 1988; Werth *et al.*, 1996b). Both tests are summarized in Fig. 1. Prolonged sleep after sleep deprivation was characterized by a monotonic decrease in SWA (Dijk *et al.*, 1990). Suppression of SWA during the initial part of time asleep without disturbing sleep caused a rebound in SWA during subsequent sleep (Dijk *et al.*, 1987b), albeit without affecting spontaneous sleep duration (Dijk and Beersma, 1989). This finding indicates that EEG SWA reflects the rate of decay of the regulating variable S, rather than S itself, as was postulated



**Figure 1.** Quantitative tests of model predictions (blue solid lines, open circles) of prior sleep and wake effects on initial non-rapid eye movement (NREM) slow wave activity (SWA). Red circles: mean normalized SWA  $\pm$  standard error of the mean (SEM) in the first half-hour of naps at different times of day (modified after Dijk *et al.*, 1987a). Red triangles: mean normalized SWA ( $\pm$  SEM) in the first half-hour of sleep at 24 h subsequent to a ca. 2-h nap at 10:00 hours and at 18:00 hours (modified after Daan *et al.*, 1988).

originally. In a further elaborated version of the model addressing changes within NREM sleep episodes, a close fit was obtained between simulated and empirical data (Achermann *et al.*, 1993).

Fewer studies have focused on the role of C in sleep timing in humans. Exposure to bright light in the morning, advancing the circadian oscillation, also advances awakening on the next day without affecting NREM sleep SWA (Dijk *et al.*, 1989). Readiness for nocturnal sleep is initiated by the circadian signal of melatonin release in the evening, which induces distal vasodilatation, enhancing the probability of sleep onset (Kräuchi and Wirz-Justice, 2001).

In animals, the rest–activity rhythm is a traditional marker of the circadian pacemaker (C). In the two-process model it is implicitly a marker of both C and S. The relationship of behaviour to S has proved important for studying sleep-like phenomena in species in which conventional electrophysiological markers are not available (e.g. invertebrates). Monitoring motor activity in the cockroach and scorpion after keeping these animals awake enhanced subsequent periods of immobility (Tobler, 1983; Tobler and Stalder, 1988). Also, brief arousals in the rat as a behavioural variable showed a negative correlation with EEG SWA (Franken *et al.*, 1991a). The use of behavioural markers of ‘sleep homeostasis’ has opened up the field of sleep research to invertebrates, and in particular to *Drosophila* and its mutants (reviewed in Faville *et al.*, 2015; Huber *et al.*, 2004). Such markers, however, preclude the assessment of an intensity dimension to sleep as in the mammalian SWA (in contrast to Faville *et al.*, 2015).

## NEW COMPLEXITIES NOT INCORPORATED INTO THE ORIGINAL MODEL

### Brain differentiation

An important development in the last two decades has been the recognition that sleep homeostasis is not only a global brain phenomenon, but also shows regional, use-dependent aspects (Krueger and Obal, 1993). Prolonged selective unihemispheric activation of the cortex during wake increased slow waves over the activated area during subsequent sleep (Kattler *et al.*, 1994). This finding was confirmed and extended using high-density EEG recordings (Huber *et al.*, 2004; Krueger and Tononi, 2011).

Regional differences in the sleep EEG have implications for the two-process model. There is a distinct antero-posterior power gradient in the NREM sleep EEG with a predominance of low-frequency power in frontal derivations (Werth *et al.*, 1996a). Analysis of the parameters of Process S have confirmed topographic differences (Rusterholz and Achermann, 2011; Zavada *et al.*, 2009). Both the increase and decrease of S are slowest in the fronto-central area. This finding may reflect the functional specialization of cortical areas (e.g. involvement of the frontal cortex in cognitive functions; reviewed in Ferrara and De Gennaro, 2011). In the context of the two-process model, such regional differences

imply that quantitative SWA-based predictions are dependent upon which brain area is chosen to estimate the time-course of S. There is currently no scientific basis for this choice.

The two-process model has strong functional implications, but it was formulated without specifying the function of sleep, except that sleep must subserve long-term maintenance of cerebral integrity. Tononi and Cirelli have proposed the synaptic homeostasis hypothesis of sleep regulation (Tononi and Cirelli, 2006, 2014). The main tenet is that synaptic and cellular processes that had been challenged during waking are re-established during sleep. Sleep is viewed as a price the brain pays for plasticity. Addressing the same issue, Frank and Cantera argue that not only sleep homeostasis but also the output of the biological clock influences brain plasticity (Frank and Cantera, 2014). Their state-clock model has received recent support from a human forced desynchrony study showing that SWA, the major index of sleep homeostasis, undergoes marked circadian variation in central and posterior cortical regions (Lazar *et al.*, 2015). Homeostatic and circadian factors seem to make synergistic contributions to synaptic plasticity.

### Continuous interaction of S and C

In the classical two-process model the sleep homeostat and the circadian process interact only at discrete events. Sleep occurs when the homeostatic Process S approaches the upper threshold of C, and waking is triggered when Process S reaches the lower threshold. This happens twice per circadian cycle under standard (monophasic) conditions, but more often with polyphasic sleep, as occurs in humans under continuous bedrest. A continuous interaction between the two processes was envisioned from the first outline of the model. The propensity for sleep or wake was assumed to depend upon the distance of S to the upper or lower thresholds of C (Borbély, 1982). Sleep propensity corresponds, therefore, to the difference between S and C. However, Process S in the model did not influence functioning of the circadian clock and sleep homeostatic processes did not change with circadian phase.

Growing evidence suggests such mutual influence, and several additive and interactive models have been proposed (Achermann, 1999; Achermann and Borbély, 2003; Dijk and Archer, 2010; Dijk and Edgar, 1999; Folkard *et al.*, 1999; Jewett and Kronauer, 1999). In particular, human data obtained in the forced desynchrony protocol have been interpreted as a non-additive interaction between sleep homeostatic mechanisms and the circadian clock (Boivin *et al.*, 1997; Dijk and Czeisler, 1995; Dijk *et al.*, 1992; Wyatt *et al.*, 1999). In this protocol, the circadian amplitude of many neurobehavioural functions was modulated by homeostatic sleep pressure. In general, a lower circadian amplitude was found when sleep pressure was high, and amplitude increased when sleep pressure was reduced. Together, the data indicate that sleep pressure affects the influence of the clock on behaviour and physiology.

In the opposite direction, the circadian clock may also influence the rise and fall of Process S. The circadian phase, where prolonged waking occurs (either spontaneous or induced), modifies the level of subsequent slow wave activity in the NREM sleep EEG (Vyazovskiy *et al.*, 2007; Werth *et al.*, 1996b). The success of predicting the level of S in a model of homeostatic sleep regulation depends upon the time of day (Deboer, 2009; Franken *et al.*, 1991b; Vyazovskiy *et al.*, 2007; Werth *et al.*, 1996b). The level of S or EEG SWA may therefore not be simply a function of previous wake duration, but may also depend upon the time of day that the waking occurs. Whether this is caused by an influence of time of day on the expression of slow waves, dissociating process S from slow wave expression or by a circadian modulation in the quality of sleep and waking is still unclear (Deboer, 2015; Lazar *et al.*, 2015).

The evidence for a role of clock genes in sleep homeostasis, reviewed elsewhere (Deboer, 2007; Franken, 2013), indicates that sleep homeostasis and circadian regulation also interact on the molecular/genetic level. Knocking out one or more of the five to seven main clock genes (*Period Per1-3*, *Clock*, *Bmal1*, *Cryptochrome Cry1* and *Cry2*, *Npas2*) not only changed or disabled the circadian clock, but also modified sleep homeostatic markers in these animal models. *Cry1-Cry2* double knockout mice show increased NREM sleep time, sleep consolidation and EEG SWA, all signs of high NREM sleep pressure. *Clock* mutants, in contrast, sleep less than wild-type control animals. *Bmal1* and *Npas2* knockout mice show altered EEG SWA and altered responses to sleep deprivation. Clock gene expression in the brain, particularly in the cerebral cortex, depends significantly upon prior sleep–wake history. Sleep deprivation during the light phase induces a duration-dependent increase in *Per2* mRNA in the mouse brain, which returns to baseline levels within 2 h of recovery sleep. These changes vary with time of day, so both factors are involved. In contrast, within the SCN, *Per2* RNA and PER2 protein expression (Curie *et al.*, 2015) remains unaffected by preceding sleep–wake history. There is no evidence that core molecular clock mechanisms in the SCN are influenced by modulations in sleep and waking.

### Interactions of SCN activity and sleep pressure

To gain more direct insight into the interaction of the circadian pacemaker and sleep, the EEG and electromyogram (EMG) have been recorded simultaneously with SCN electrical neuronal activity *in vivo* in freely moving rats. SCN neuronal activity is high during the subjective day and low during the subjective night, irrespective of whether an animal is diurnal or nocturnal (Sato and Kawamura, 1984; Vansteensel *et al.*, 2008). In such simultaneous recordings it was shown that SCN neuronal activity increases during waking and REM sleep and decreases during NREM sleep, in addition to the well-known circadian changes. These ultradian modulations correlate with the level of SWA in the different sleep states. In addition, careful slow wave deprivation during NREM sleep,

reducing SWA but still allowing the animals to sleep, results in increased SCN neuronal activity (Deboer *et al.*, 2003). Increasing SWA by sleep deprivation leads to a decrease in SCN neuronal activity in NREM sleep and REM sleep (Deboer *et al.*, 2007).

This relationship between SCN neuronal activity and sleep pressure in rats has also been found in humans. The fMRI BOLD signal, recorded during a psychomotor vigilance task, declined in the suprachiasmatic area when sleep pressure was increased (Schmidt *et al.*, 2009). The change in BOLD signal correlated with SWA in the first NREM sleep episode. Thus, the circadian pacemaker appears to obtain feedback from the status of the sleep homeostat, such that increased sleep pressure reduces circadian amplitude.


### Sleep pressure changes and clock functioning

Interactions between sleep pressure changes and circadian clock functioning have been observed in rodents. In mice and hamsters, sleep deprivation attenuates phase shifts induced by light (Challet *et al.*, 2001; Mistlberger *et al.*, 1997; Van Diepen *et al.*, 2014), suggesting that increased sleep pressure reduces circadian clock response to the zeitgeber. Similarly, in humans, attenuated phase advances were found after sleep restriction (Burgess, 2010).

Light shifts the clock in the SCN via retinal ganglion cells and a monosynaptic pathway, the retinohypothalamic tract. Following activation by light, glutamate is released at the nerve terminals (Ding *et al.*, 1994; Johnson *et al.*, 1988), leading to an increase in SCN neuronal activity (Meijer *et al.*, 1998; Van Diepen *et al.*, 2013). Application of glutamate to the SCN mimics the effect of light (Ding *et al.*, 1994). Sleep deprivation may reduce the phase-shifting capacity of light by diminishing the strength of the photic signal reaching the SCN, possibly through blocking glutamate release.

Adenosine is thought to be involved in the homeostatic regulation of sleep (Landolt, 2008), and its extracellular level in the brain increases in the course of prolonged waking (Porkka-Heiskanen *et al.*, 1997). In various brain regions, the stimulation of presynaptic A1 adenosine receptors depresses glutamate release and reduces the amplitude of excitatory postsynaptic currents (Barrie and Nicholls, 1993; Dolphin and Prestwich, 1985; Oliet and Poulain, 1999). This may be the mechanism whereby increased adenosine levels after sleep deprivation influence the light responsiveness of the circadian clock. Indeed, after a 6-h sleep deprivation ending at CT14, the light response in the SCN was reduced compared to the control; systemic injection of caffeine restored this attenuation of the SCN light response almost completely (Van Diepen *et al.*, 2014), supporting the interactive role of adenosine and glutamate. The reduced response to light of SCN neuronal activity after sleep deprivation provides evidence that clock functioning may be modified by sleep homeostatic pressure.

Figure 2 summarizes some changes in clock output and functioning under the influence of increased sleep pressure.



**Figure 2.** Selected effects of sleep deprivation on circadian clock modulated brain and behavioural functions. Sleep deprivation reduces light-induced behavioural phase shifts (Challet *et al.*, 2001; Mistlberger *et al.*, 1997; Van Diepen *et al.*, 2014) and the amplitude of the circadian modulation of SCN neuronal activity (Deboer *et al.*, 2007). In humans it reduces the fMRI BOLD response during a psychomotor vigilance task (Schmidt *et al.*, 2009). In rodents it reduces the light-induced increase in firing rate in SCN neurones (Van Diepen *et al.*, 2014).

Thus, it can be concluded that a continuous exchange of information between the two processes takes place. The homeostatic and circadian processes are not simply independent or additive, but probably subject to more complex interactions. Future modelling efforts should consider the strength of clock output as a function of sleep homeostatic pressure.

### Sleep–wake timing controlled by a circadian oscillator outside the SCN

The original view of Process S as a global cortical process has been modified to include regional and use-dependent facets. Similarly, the traditional view of Process C as a hierarchical circadian system under the strict control of the SCN has been expanded by the recognition of peripheral circadian oscillators that exhibit a high degree of

independence and flexibility and may respond to physiological and behavioural concomitants of sleep and waking. Thus, as had been shown for sleep homeostasis, a global (systemic) and a local regulation can be distinguished also for circadian rhythms (Asher and Schibler, 2011). These new insights have been obtained in the domain of chronobiology and their repercussions with respect to sleep have not yet been examined closely.

There is accumulating evidence that circadian rest–activity cycles generated outside the SCN can emerge in different organisms under non-standard (laboratory) conditions. In several field studies, rodents that are typically nocturnal in the laboratory may become diurnal under naturalistic conditions: spiny mice (Levy *et al.*, 2007), Syrian hamster (Gattermann *et al.*, 2008), house mouse (Daan *et al.*, 2011) and Tuco-tuco (Tomotani *et al.*, 2012). The degu, a species considered diurnal, becomes nocturnal when given a running wheel in the laboratory, while the circadian pacemaker, as judged from its light response curve, remains at the original phase (Kas and Edgar, 2000). Circadian rhythmicity in rest–activity can even be expressed in the absence of the SCN, as found in SCN-lesioned rats under temporal food restriction, driven apparently by a separate oscillator (Stephan *et al.*, 1979). Rats and mice given methamphetamine in their drinking water show pronounced circadian rest–activity rhythms that run typically with longer periods no longer entrained to about 24 h by the SCN, and even persist in the absence of the SCN (Honma and Honma, 2009). This methamphetamine-sensitive rhythm was shown to be independent of the canonical SCN clock genes *cry1* and *cry2* (Honma *et al.*, 2008) or *per1*, *per2*, *bmal1*, *npas2*, *clock* and *Ck1ε* (Mohawk *et al.*, 2009). The nature of this circadian oscillator outside the SCN is still enigmatic, but it may well be identical with the food entrainable oscillator, as the rhythm of methamphetamine-treated, SCN-lesioned rats entrains to restricted daily feeding (Honma *et al.*, 1989). Thus, it has become likely that the overt rest–activity rhythm is generated by a circadian oscillator outside the SCN that, under standard conditions, maintains a rather fixed-phase relationship with the SCN and thereby with the light–dark cycle. Recently, a dopaminergic ultradian oscillator has been identified in the mouse which normally cycles in harmony with the circadian clock, but can desynchronize when dopaminergic tone is elevated (Blum *et al.*, 2014). This can give rise to aberrant behavioural oscillations, and may account for the methamphetamine-induced cyclicality. In this view, the role of the SCN is not so much the generation of activity rhythms, but rather the maintenance of temporal integrity of circadian physiology and behaviour with external day and night. The loss of control over sleep–wake oscillations in human internal desynchronization has been explained by a reduced amplitude of Process C in the absence of resonance with any zeitgeber (Daan *et al.*, 1984), but this does not disprove that sleep and wake could themselves be generated by an as yet unknown oscillator outside the SCN.

### A role for metabolism in the control of rest–activity timing

As reviewed above, the circadian rhythm of activity and rest can adopt different phase relationships with the pacemaker. Animals that are diurnal in the field can become nocturnal in the laboratory. Mice that are nocturnal in the laboratory become diurnal when they do not receive their food *ad libitum*, as is standard in the laboratory, but have to work to obtain it, as is usual in nature (Hut *et al.*, 2011). The gradual shift of activity from night to day is followed eventually by the occurrence of torpor (body temperature approximately 23 °C) in the late night. Both low temperature and the working-for-food protocol shift mouse activity from night into day, even though the SCN retains its original phase relationship to the light–dark cycle: a rigid clock driving flexible rhythms (Van der Vinne *et al.*, 2014). These findings strongly support a role for energy metabolism in determining the phase of the rest–activity rhythm relative to the SCN. The emergence of activity rhythms in SCN-lesioned rodents under restricted feeding (Stephan *et al.*, 1979) indicates that the amplitude of the underlying oscillator is also affected by energy metabolism.

A further potential link between human rest–activity cycles and energy metabolism is present in the data obtained by Aschoff and coworkers in subjects under isolation from time cues. During internal desynchronization, the time between meals varied in proportion to the duration of wakefulness (varying from 12 to 42 h) (Aschoff *et al.*, 1986). Indeed, caloric intake per cycle (Green *et al.*, 1987) and activity per cycle (Aschoff, 1993) remained the same, despite very large changes in cycle duration. Aschoff deduced early on that the length of the rest–activity cycle may be associated negatively with basal metabolic rate (Aschoff, 1993). While energy metabolism was not assessed in these studies, subsequent analyses demonstrated a negative association between cycle length and core body temperature (Daan *et al.*, 2013). To the extent that the variation in body temperature reflects the variation in heat production (metabolic rate), the results are consistent with the proposition that the spontaneous frequency of the human sleep–wake oscillator is associated with metabolic rate, as suggested on the basis of the proportionality of meal frequency and sleep–wake frequency. The original S–C model may still provide a framework for the much-needed analysis of the role of metabolism in the temporal organization of sleep and wake, and its integration with sleep homeostasis. However, this should account for the role of an oscillator outside the SCN, and outside the realm of the canonical transcriptional oscillator in the SCN. The big unanswered experimental question here is how the rise of S and the spontaneous timing of sleep and wake depend upon metabolic rate.

### Temporal segregation of metabolic processes

The field of circadian rhythms also evolved in another aspect. Gene transcription is not a prerequisite for circadian rhythms,


and non-transcriptional events are sufficient to sustain circadian rhythms in human red blood cells (O'Neill and Reddy, 2011). Cellular circadian rhythms may share a common molecular origin. The oxidation–reduction cycles of peroxiredoxin proteins has been proposed to constitute a universal marker for circadian rhythms in all domains of life (Edgar *et al.*, 2012). Metabolic cycles and circadian oscillators may be linked so inextricably that little meaningful distinction can be made between them. One of the central functional aspects is the temporal segregation of mutually antagonistic metabolic processes.

In the framework of chronobiology, sleep is viewed increasingly in the context of metabolism. Metabolic processes are modulated by time of day and via synchronization of the circadian pacemaker to the light–dark cycle; thereby, predominantly anabolic and predominantly catabolic processes in peripheral tissues are segregated. Thus, in the wake-feeding episode glycogen and cholesterol synthesis are promoted, whereas in the sleep-fasting episode gluconeogenesis and glycogenolysis prevail (Bass and Takahashi, 2010). In terms of hormone secretion, release of glucocorticoids and insulin dominates in the former, with release of melatonin, growth hormone and leptin in the latter. It is noteworthy that the gut microbiome is highly dynamic, exhibiting daily cyclic fluctuations which have repercussions on host metabolism and provide evidence for a cross-regulation of prokaryotic and eukaryotic circadian rhythms (Thaiss *et al.*, 2014; Zarrinpar *et al.*, 2014). We could view the microbiome as forming part of our internal environment.

Contemporary chronobiology appears to revive concepts advanced more than 80 years ago. Walter Rudolf Hess (Hess, 1932) advocated two functional phases of the autonomic nervous system: the ergotropic phase is characterized by the predominance of the sympathetic nervous system, enabling interaction with the external environment such as in foraging, social interaction and predator avoidance. The trophotropic phase with the predominance of the parasympathetic system is the opposite – with an energy-preserving, stress-preventing and restorative function (Fig. 3). According to Hess, sleep is a vegetative process by which the autonomic nervous system regulates activity of higher cerebral functions.

The special mechanisms which bring about repair during sleep are hidden in the tissues. They have not yet been fully explained; their existence is only deduced from their effects; yet they lie at the heart of the problem of sleep, and the resting of the sense-organs, muscles, and psychic functions are only accessory factors facilitating restoration within the tissues (Hess, 1932).

In other words, by enforcing quiescence and fasting, sleep ensures optimal physiological and behavioural conditions for restorative peripheral metabolic processes to occur. In this respect, sleep is an adjunct of the 24-h cycle of metabolism. If sleep does not occur within the appropriate phase of the cycle, by being either too short or too long, the risk of type 2



**Figure 3.** Functional antagonism within the circadian cycle. Within the 24-h cycle, Hess distinguished a trophotropic phase with parasympathetic predominance characterized by resting, fasting and anabolism, and an ergotropic phase with sympathetic predominance characterized by motor activity, feeding, drinking and catabolism. Hess’s concept has been revived in contemporary rhythm research by the circadian clock’s partitioning of behavioural and metabolic processes according to the time of day.

diabetes increases (Cappuccio *et al.*, 2010). Delaying the habitual sleep phase reduces insulin sensitivity and enhances a marker for inflammation, presumably by exposing peripheral organs to nutrients during the habitual overnight fast period (Leproult *et al.*, 2014).

Sleep has been shown to be an important factor for maintaining rhythmicity of circadian transcripts in the periphery.


When sleep is phase-shifted with respect to the clock, the majority of circadian transcripts become arrhythmic (Archer *et al.*, 2014). Thus, sleep occurring at its habitual circadian phase engenders ‘resonance’ between central and peripheral cues.

**IMPACT OF THE TWO-PROCESS MODEL ON CLINICAL RESEARCH**


**Sleep deprivation**

Formulation of the two-process model provided an immediate tool for depression research. In the 1970s, Pflug and Tölle had systematically investigated the remarkably rapid antidepressant effect of one night’s sleep deprivation in affective disorders (Pflug and Tölle, 1971). As documented in thousands of patients, the depression usually lifts towards the morning after a complete night awake, and often returns following the recovery night sleep. This clinical observation lent itself to the question as to whether such a rapid improvement and rapid relapse could be related to and modelled by abnormalities in Process S (Borbély and Wirz-Justice, 1982) (Fig. 4). The S-deficiency hypothesis of depression did not specify a neurochemical/molecular mechanism; its merit lies in the simple proposal that a deficient S-process is normalized after sleep deprivation. A second


**(a) Markers for model parameters**



**(b) Disturbances in depression and model predictions for treatment**



**Figure 4.** Schematic of the two-process model and its application to depression. Normal situation with markers for model parameters (left) and putative pathological changes and their remedies (right). The skewed sinusoidal function of C corresponds to the quantitative version of the model (Daan *et al.*, 1984) and does not represent the time-course of the markers.



**Figure 5.** Timing of sleep and mood – the internal coincidence model of sleep deprivation (Wehr and Wirz-Justice, 1981).

‘internal coincidence’ model additionally incorporated Process C: the appropriate timing of sleep with respect to the internal clock was crucial for stable mood (Wehr and Wirz-Justice, 1981) (Fig. 5). This latter model developed from clinical manipulations of sleep length and timing in depressive patients – partial sleep deprivation in the second half of the night was as efficacious as total sleep deprivation. Even phase-advancing the sleep–wake cycle (without deprivation of sleep) was antidepressant and, although it took longer for the effect to develop, it also lasted longer (Wirz-Justice and Van Den Hoofdakker, 1999) (Fig. 5). There was an initial flurry of sleep EEG studies after publication of the hypothesis, but only recently has there been a rekindling of interest in the mechanisms underlying clinical sleep deprivation using new imaging technologies, more stringent protocols and carefully diagnosed patient groups, following markers of both processes S and C. A very concise, extensive new review of sleep deprivation in depression covers all aspects of putative mechanisms from imaging to synaptic plasticity (Dallaspezia and Benedetti, 2015).

### Seasonal affective disorder

The most consistent and careful studies based on the model, using both forced desynchrony and constant routine protocols, have been carried out in patients with winter seasonal affective disorder (SAD). Remarkably few differences between SAD and controls in terms of circadian period, phase or sleep SWA were observed (Brunner *et al.*, 1996; Koorengel *et al.*, 2002a,b; Wirz-Justice *et al.*, 1995). Studies of classical photoreceptor function (ERG, EOG, dark adaptation) predominantly show normal retinal parameters in SAD in summer and subsensitivity to environmental light in winter; however, the data are not quite consistent (Danilenko and Levitan, 2012). More specifically, SAD patients may have a decreased retinal sensitivity related to melanopsin gene variations in the non-image-forming light-input pathway as measured by the post-illumination pupil response (Roeklein *et al.*, 2013). Functional magnetic resonance imaging

(fMRI) has revealed abnormal hypothalamic responses to light in SAD (Vandewalle *et al.*, 2011).

We still do not know why SAD patients are vulnerable to diminishing photoperiod in winter (Danilenko and Levitan, 2012). However, the model provided a conceptual framework to test hypotheses of SAD pathophysiology rigorously and, most importantly, to investigate the clinical effects of the zeitgeber light.

### Light as therapy

Light for SAD was the first treatment in psychiatry based on neurobiological research – the simulation of a spring photoperiod to counteract the long winter night (Rosenthal *et al.*, 1984). Later, it was found that an early dawn signal was sufficient for clinical response in SAD (Terman and Terman, 2006): part of this improvement is related to the phase advance induced by morning light, but also to direct effects of light on mood, probably via the same serotonergic systems as found responding to classical selective serotonin reuptake inhibitors (SSRIs) (Spindelegger *et al.*, 2012). Light does not modify sleep EEG spectra in SAD (Brunner *et al.*, 1996). According to the model (Fig. 4), light acts on Process C to shift phase and increase amplitude, and by increasing zeitgeber strength it stabilizes entrainment. The direct neurochemical effects of light may explain why it is effective at different times of day, the circadian modulation being relevant for better morning response.

There is an impressive literature on light’s multiple actions on the brain; in particular, the non-visual effects of light transduced by wavelength-dependent novel melanopsin-containing photoreceptors (ipRGCs) (Gaggioni *et al.*, 2014). For example, an aberrant light cycle that neither changed the amount and architecture of sleep nor caused changes in the circadian timing system directly regulated mood-related behaviours and cognitive functions in mice via the ipRGCs (Legates *et al.*, 2012).

### Major depression

Even though the two-process model was applied initially to major depression, there have been few studies in these patients, with results ranging from the early finding of low NREM EEG delta power (Borbély *et al.*, 1984) to higher SWA in young depressed women (Frey *et al.*, 2012), who show a stronger response to sleep deprivation than controls, particularly in frontal brain regions. High-density EEG demonstrated higher SWA primarily in prefrontal channels only in depressed women (Plante *et al.*, 2012). In contrast, another study found lower SWA in outpatients with depression, particularly men aged 20–30 years (Brower *et al.*, 2011). We are not aware of any stringent investigations of sleep and circadian rhythms in major non-seasonal depression or bipolar disorder under constant routine or forced desynchrony protocols.

Evidence for the tenets of the two-process model has been found in the antidepressant effects of selective slow wave

sleep deprivation (Landsness *et al.*, 2011). The reduction in depressive symptoms correlated with the overnight dissipation of fronto-central SWA during baseline sleep, and the rebound in right frontal all-night SWA during recovery sleep.

Measures of cortical responses evoked by transcranial magnetic stimulation (TMS) have been correlated with synaptic strength in rodents and change with sleep homeostatic pressure in humans. TMS/EEG in the prefrontal cortex in bipolar depression reveal that cortical excitability, putatively a direct measure of Process S, increases as a function of time awake, more in clinical responders than in non-responders (Canali *et al.*, 2014). Dallaspezia and Benedetti (2015) conclude in their review: 'Moreover, considering that cortical excitability has been proposed as a correlate of process S, these findings then confirm hypotheses of process S increase as a core mechanism of action of antidepressant SD'.

A striking development in the clinical domain has been a new generation of treatments combining manipulations of Processes S and C: sleep deprivation with light therapy – yielding a fast and long-lasting response (e.g. Benedetti *et al.*, 2014; Martiny *et al.*, 2012; Wu *et al.*, 2009), some even including a sleep phase advance (Echizenya *et al.*, 2013; Sahlem *et al.*, 2014). Thus, the two components of the model may not be understood clearly, but the treatments arising therefrom are already established (Wirz-Justice *et al.*, 2013).

### Beyond depression

Many abnormal sleep patterns in psychiatry and other medical disciplines can be understood more clearly when considering the contributions of S and C in order to specify the disturbance. As an example, arrhythmic rest–activity cycles in a schizophrenic patient treated with the sedating antipsychotic haloperidol led to measurement of melatonin rhythms as a circadian marker (Wirz-Justice *et al.*, 1997). Surprisingly, the melatonin rhythm persisted, indicating a functioning clock. In the two-process model, the frequency of sleep–wake alternations depends upon the distance between the circadian thresholds defining sleep and wake. In simulations, reducing this interval induces polyphasic sleep (Daan *et al.*, 1984), leading to the question of whether the drug-induced sedation could have been the reason for the poor rest–activity pattern. This led to a change in medication followed by the re-emergence of a circadian pattern.

A broad field of application has been in alertness management (fatigue and performance), development (adolescence) and ageing, dissecting out reasons for individual differences (long/short sleepers, early/late chronotype, genotype). It has led in new directions, to investigations of trait-like individual responses to sleep loss or circadian misalignment, time-on-task effects, cumulative effects of sleep restriction, executive function and local versus global sleep. Although not developed initially for clinical application, the model has unravelled the two processes contributing to many

physiological, behavioural and cognitive functions. This research has yielded novel, non-pharmacological antidepressant treatments such as light therapy, sleep deprivation and sleep phase advance.

### CONCLUDING REMARKS

The two-process model continues to be conceptually useful for organizing the thinking about sleep regulation along two axes. It is still helpful for bringing the disciplines of sleep and rhythm research closer together. It also remains influential in the design of experiments and data evaluation. In medicine, disease may induce non-specific impairment of homeostatic mechanisms. Illness may affect light-orientated behaviour that has consequences for stable entrainment. Even in its classical form, the two-process model provides a simple construct to differentiate sleep disturbances appropriately and thereby initiate specific treatments.

The detection of local, use-dependent changes of the sleep EEG has provided regional and functional specificity of the effect of waking on sleep. Recent neurophysiological and morphological studies have focused upon local sleep-related changes at the synaptic level and their functional implications. It is important to be aware that plasticity in the brain is modified by both homeostatic and circadian factors. Some of the detrimental effects of sleep deprivation are due to the disruption of the synergy of S and C.

Our view of the circadian system has undergone profound changes. The SCN is now viewed as orchestrating and integrating rhythms rather than simply generating and driving them. Homeostasis of (still imprecisely defined aspects of) brain function may be achieved through periodic sleep–wake alternation, which itself is generated by either an underlying neuronal oscillator outside the SCN or by a behavioural relaxation oscillator as conceived in Process S. The verdict on this issue is still pending. Slow advancing or delaying transients of experimentally shifted human sleep–wake cycles towards the unshifted circadian melatonin rhythm in temporal isolation (Hashimoto *et al.*, 2004) are consistent with an underlying sleep–wake oscillator. Other researchers have conceptualized a relaxation-type sleep–wake switch involving both orexinergic wake-promoting and homeostatic sleep-promoting drives (Phillips *et al.*, 2011). When the orexinergic wake promotion is under circadian control and reflects the daily rise in Process C, this is fully compatible with the two-process model.

Sleep has a large effect on peripheral circadian oscillators due to such factors as the absence of locomotion, reduced body temperature, increased growth hormone and reduced glucocorticoid secretion. If sleep occurs at the optimal circadian phase, it reinforces the temporal segregation of 'sleep/fast-type' metabolism from 'wake/feed-type' metabolism. In addition to a need-related facet that depends on previous history, sleep ensures an optimal internal environment for processes occurring during the circadian resting phase. In an early paper, the terms 'recovery sleep' and

'inactivity sleep' were used to refer to the two facets of sleep (Borbély, 1980). It was argued that the latter type of sleep ensures the immobility of the animal during specific parts of the 24-h cycle, thereby minimizing energy expenditure and possibly reducing the risk of predation. Three decades later, we emphasize the metabolic benefits of inactivity. The possibility of modulating sleep intensity in addition to duration contributes functionally to optimal flexibility in the circadian timing of sleep.

## REFERENCES

- Achermann, P. Technical note. A problem with identifying nonlinear interactions of circadian and homeostatic processes. *J. Biol. Rhythms*, 1999, 14: 602–603.
- Achermann, P. and Borbély, A. A. Mathematical models of sleep regulation. *Front. Biosci. (Landmark edn.)*, 2003, 8: S683–S693.
- Achermann, P., Dijk, D. J., Brunner, D. P. and Borbély, A. A. A model of human sleep homeostasis based on EEG slow-wave activity – quantitative comparison of data and simulations. *Brain Res. Bull.*, 1993, 31: 97–113.
- Archer, S. N., Laing, E. E., Moeller-Levet, C. S. *et al.* Mistimed sleep disrupts circadian regulation of the human transcriptome. *Proc. Natl Acad. Sci. USA*, 2014, 111: E682–E691.
- Aschoff, J. On the relationship between motor activity and the sleep–wake cycle in humans during temporal isolation. *J. Biol. Rhythms*, 1993, 8: 33–46.
- Aschoff, J., Von Goetz, C., Wildgruber, C. and Wever, R. A. Meal timing in humans during isolation without time cues. *J. Biol. Rhythms*, 1986, 1: 151–162.
- Asher, G. and Schibler, U. Crosstalk between components of circadian and metabolic cycles in mammals. *Cell Metab.*, 2011, 13: 125–137.
- Barrie, A. P. and Nicholls, D. G. Adenosine A1 receptor inhibition of glutamate exocytosis and protein kinase C-mediated decoupling. *J. Neurochem.*, 1993, 60: 1081–1086.
- Bass, J. and Takahashi, J. S. Circadian integration of metabolism and energetics. *Science*, 2010, 330: 1349–1354.
- Benedetti, F., Riccaboni, R., Locatelli, C., Poletti, S., Dallaspezia, S. and Colombo, C. Rapid treatment response of suicidal symptoms to lithium, sleep deprivation, and light therapy (Chronotherapeutics) in drug-resistant bipolar depression. *J. Clin. Psychiatry*, 2014, 75: 133–140.
- Blum, I. D., Zhu, L., Moquin, L. *et al.* A highly-tunable dopaminergic oscillator generates ultradian rhythms of behavioral arousal. *eLife*, 2014, 3: e05105.
- Boivin, D. B., Czeisler, C. A., Dijk, D. J. *et al.* Complex interaction of the sleep–wake cycle and circadian phase modulates mood in healthy subjects. *Arch. Gen. Psychiatry*, 1997, 54: 145–152.
- Borbély, A. A. Sleep: circadian rhythm versus recovery process. In: M. Koukkou, D. Lehmann and J. Angst (Eds) *Functional States of the Brain: Their Determinants*. Elsevier, Amsterdam, 1980: 151–161.
- Borbély, A. A. A two-process model of sleep regulation. *Hum. Neurobiol.*, 1982, 1: 195–204.
- Borbély, A. A. and Achermann, P. Sleep homeostasis and models of sleep regulation. *J. Biol. Rhythms*, 1999, 14: 557–568.
- Borbély, A. A. and Wirz-Justice, A. Sleep, sleep deprivation and depression. A hypothesis derived from a model of sleep regulation. *Hum. Neurobiol.*, 1982, 1: 205–210.
- Borbély, A. A., Tobler, I., Loeferle, M. *et al.* All-night spectral analysis of the sleep EEG in untreated depressives and normal controls. *Psychiatry Res.*, 1984, 12: 27–33.
- Brower, K. J., Hoffmann, R., Conroy, D. A., Arndt, J. T. and Armitage, R. Sleep homeostasis in alcohol-dependent, depressed and healthy control men. *Eur. Arch. Psychiatry Clin. Neurosci.*, 2011, 261: 559–566.
- Brunner, D. P., Kräuchi, K., Dijk, D. J., Leonhardt, G., Hang, H. J. and Wirz-Justice, A. Sleep electroencephalogram in seasonal affective disorder and in control women: effects of midday light treatment and sleep deprivation. *Biol. Psychiatry*, 1996, 40: 485–496.
- Burgess, H. J. Partial sleep deprivation reduces phase advances to light in humans. *J. Biol. Rhythms*, 2010, 25: 460–468.
- Canali, P., Papa, G. S., Casali, A. G. *et al.* Changes of cortical excitability as markers of antidepressant response in bipolar depression: preliminary data obtained by combining transcranial magnetic stimulation (TMS) and electroencephalography (EEG). *Bipolar Disord.*, 2014, 16: 809–819.
- Cappuccio, F. P., D'elia, L., Strazzullo, P. and Miller, M. A. Quantity and quality of sleep and incidence of type 2 diabetes – a systematic review and meta-analysis. *Diabetes Care*, 2010, 33: 414–420.
- Challet, E., Turek, F. W., Laute, M. A. and Van Reeth, O. Sleep deprivation decreases phase-shift responses of circadian rhythms to light in the mouse: role of serotonergic and metabolic signals. *Brain Res.*, 2001, 909: 81–91.
- Curie, T., Maret, S., Emenegger, Y. and Franken, P. *In vivo* imaging of the central and peripheral effects of sleep deprivation and suprachiasmatic nuclei lesion on PERIOD-2 protein in mice. *Sleep*, 2015, 38: 1381–1394.
- Daan, S., Beersma, D. G. M. and Borbély, A. A. Timing of human sleep: recovery process gated by a circadian pacemaker. *Am. J. Physiol.*, 1984, 246: R161–R178.
- Daan, S., Beersma, D. G. M., Dijk, D. J., Åkerstedt, T. and Gillberg, M. Kinetics of an hourglass component involved in the regulation of human sleep and wakefulness. In: W. T. J. M. Hekken, G. A. Kerkhof and W. J. Rietveld (Eds) *Advances in the Biosciences*, vol. 73. Trends in Chronobiology; XVIIIth Conference of the International Society for Chronobiology in Conjunction with the IIIrd Annual Meeting of the European Society for Chronobiology, Leiden, Netherlands, 12–17 July 1987. XIII+386 pp. Pergamon Press, Oxford; New York. Illus. 1988: 183–194.
- Daan, S., Spoelstra, K., Albrecht, U. *et al.* Lab mice in the field: unorthodox daily activity and effects of a dysfunctional circadian clock allele. *J. Biol. Rhythms*, 2011, 26: 118–129.
- Daan, S., Honma, S. and Honma, K. Body temperature predicts the direction of internal desynchronization in humans isolated from time cues. *J. Biol. Rhythms*, 2013, 28: 403–411.
- Dallaspezia, S. and Benedetti, F. Sleep deprivation therapy for depression. *Curr. Top. Behav. Neurosci.*, 2015, 25: 483–502.
- Danilenko, K. V. and Levitan, R. D. Seasonal affective disorder. *Handb. Clin. Neurol.*, 2012, 106: 279–289.
- Deboer, T. Technologies of sleep research. *Cell. Mol. Life Sci.*, 2007, 64: 1227–1235.
- Deboer, T. Sleep and sleep homeostasis in constant darkness in the rat. *J. Sleep Res.*, 2009, 18: 357–364.
- Deboer, T. Behavioral and electrophysiological correlates of sleep and sleep homeostasis. *Curr. Top. Behav. Neurosci.*, 2015, 25: 1–24.
- Deboer, T., Vansteensel, M. J., Detari, L. and Meijer, J. H. Sleep states alter activity of suprachiasmatic nucleus neurons. *Nat. Neurosci.*, 2003, 6: 1086–1090.
- Deboer, T., Detari, L. and Meijer, J. H. Long term effects of sleep deprivation on the mammalian circadian pacemaker. *Sleep*, 2007, 30: 257–262.
- Dijk, D. J. and Archer, S. N. PERIOD3, circadian phenotypes, and sleep homeostasis. *Sleep Med. Rev.*, 2010, 14: 151–160.
- Dijk, D. J. and Beersma, D. G. M. Effects of SWS deprivation on subsequent EEG power density and spontaneous sleep duration. *Electroencephalogr. Clin. Neurophysiol.*, 1989, 72: 312–320.

- Dijk, D. J. and Czeisler, C. A. Contribution of the circadian pacemaker and the sleep homeostat to sleep propensity, sleep structure, electroencephalic slow waves, and sleep spindle activity in humans. *J. Neurosci.*, 1995, 15: 3526–3538.
- Dijk, D. J. and Edgar, D. M. Circadian and homeostatic control of wakefulness and sleep. In: F. W. Turek and P. C. Zee (Eds) *Regulation of Sleep and Circadian Rhythms*. Marcel Dekker Inc, New York, Basel, 1999: 111–147.
- Dijk, D. J., Beersma, D. G. M. and Daan, S. EEG power density during nap sleep: reflection of an hourglass measuring the duration of prior wakefulness. *J. Biol. Rhythms*, 1987a, 2: 207–219.
- Dijk, D. J., Beersma, D. G. M., Daan, S., Bloem, G. and Van Den Hoofdakker, R. H. Quantitative analysis of the effects of slow wave sleep deprivation during the first 3 h of sleep on subsequent EEG power density. *Eur. Arch. Psychiatry Neurol. Sci.*, 1987b, 236: 323–328.
- Dijk, D. J., Beersma, D. G. M., Daan, S. and Lewy, A. J. Bright morning light advances the human circadian system without affecting NREM sleep homeostasis. *Am. J. Physiol.*, 1989, 256: R106–R111.
- Dijk, D. J., Brunner, D. P. and Borbély, A. A. Time course of EEG power density during long sleep in humans. *Am. J. Physiol.*, 1990, 258: R650–R661.
- Dijk, D. J., Duffy, J. F. and Czeisler, C. A. Circadian and sleep/wake dependent aspects of subjective alertness and cognitive performance. *J. Sleep Res.*, 1992, 1: 112–117.
- Ding, J. M., Chen, D., Weber, E. T., Faiman, L. E., Rea, M. A. and Gillette, M. U. Resetting the biological clock: mediation of nocturnal circadian shifts by glutamate and NO. *Science*, 1994, 266: 1713–1717.
- Dolphin, A. C. and Prestwich, S. A. Pertussis toxin reverses adenosine inhibition of neuronal glutamate release. *Nature*, 1985, 316: 148–150.
- Echizenya, M., Suda, H., Takeshima, M., Inomata, Y. and Shimizu, T. Total sleep deprivation followed by sleep phase advance and bright light therapy in drug-resistant mood disorders. *J. Affect. Disord.*, 2013, 144: 28–33.
- Edgar, R. S., Green, E. W., Zhao, Y. *et al.* Peroxiredoxins are conserved markers of circadian rhythms. *Nature*, 2012, 485: 459–465.
- Faville, R., Kottler, B., Goodhill, G. J., Shaw, P. J. and Van Swinderen, B. How deeply does your mutant sleep? Probing arousal to better understand sleep defects in *Drosophila*. *Sci. Rep.*, 2015, 5: 1–13.
- Ferrara, M. and De Gennaro, L. Going local: insights from EEG and stereo-EEG studies of the human sleep–wake cycle. *Curr. Top. Med. Chem.*, 2011, 11: 2423–2437.
- Folkard, S., Åkerstedt, T., Macdonald, I., Tucker, P. and Spencer, M. B. Beyond the three-process model of alertness: estimating phase, time on shift, and successive night effects. *J. Biol. Rhythms*, 1999, 14: 577–587.
- Frank, M. G. and Cantera, R. Sleep, clocks, and synaptic plasticity. *Trends Neurosci.*, 2014, 37: 491–501.
- Franken, P. A role for clock genes in sleep homeostasis. *Curr. Opin. Neurobiol.*, 2013, 23: 864–872.
- Franken, P., Dijk, D. J., Tobler, I. and Borbély, A. A. Sleep deprivation in rats: effects on EEG power spectra, vigilance states, and cortical temperature. *Am. J. Physiol.*, 1991a, 261: R198–R208.
- Franken, P., Tobler, I. and Borbély, A. A. Sleep homeostasis in the rat: simulation of the time course of EEG slow-wave activity. *Neurosci. Lett.*, 1991b, 130: 141–144.
- Frey, S., Birchler-Pedross, A., Hofstetter, M. *et al.* Young women with major depression live on higher homeostatic sleep pressure than healthy controls. *Chronobiol. Int.*, 2012, 29: 278–294.
- Gaggioni, G., Maquet, P., Schmidt, C., Dijk, D. J. and Vandewalle, G. Neuroimaging, cognition, light and circadian rhythms. *Front. Syst. Neurosci.*, 2014, 8: 126.
- Gattermann, R., Johnston, R. E., Yigit, N. *et al.* Golden hamsters are nocturnal in captivity but diurnal in nature. *Biol. Lett.*, 2008, 4: 253–255.
- Green, J., Pollak, C. P. and Smith, G. P. Meal size and intermeal interval in human subjects in time isolation. *Physiol. Behav.*, 1987, 41: 141–147.
- Hashimoto, S., Nakamura, K., Honma, S. and Honma, K. Non-photic entrainment of human rest–activity cycle independent of circadian pacemaker. *Sleep Biol. Rhythms*, 2004, 2: 29–36.
- Hess, W. R. The autonomic nervous system. *Lancet*, 1932, 2: 1259–1261.
- Honma, K. and Honma, S. The SCN-independent clocks, methamphetamine and food restriction. *Eur. J. Neurosci.*, 2009, 30: 1707–1717.
- Honma, S., Honma, K. and Hiroshige, T. Methamphetamine induced locomotor rhythm entrains to restricted daily feeding in SCN lesioned rats. *Physiol. Behav.*, 1989, 45: 1057–1065.
- Honma, S., Yasuda, T., Yasui, A., Van Der Horst, G. T. J. and Honma, K. Circadian behavioral rhythms in *Cry1/Cry2* double-deficient mice induced by methamphetamine. *J. Biol. Rhythms*, 2008, 23: 91–94.
- Huber, R., Hill, S. L., Holladay, C., Biesiadecki, M. and Tononi, G. Sleep homeostasis in *Drosophila melanogaster*. *Sleep*, 2004, 27: 628–639.
- Hut, R. A., Pilonz, V., Boerema, A. S., Strijkstra, A. M. and Daan, S. Working for food shifts nocturnal mouse activity into the day. *PLoS One*, 2011, 6: 1–6.
- Jewett, M. E. and Kronauer, R. E. Interactive mathematical models of subjective alertness and cognitive throughput in humans. *J. Biol. Rhythms*, 1999, 14: 588–597.
- Johnson, R. F., Moore, R. Y. and Morin, L. P. Loss of entrainment and anatomical plasticity after lesions of the hamster retinohypothalamic tract. *Brain Res.*, 1988, 460: 297–313.
- Kas, M. J. H. and Edgar, D. M. Photic phase response curve in *Octodon degus*: assessment as a function of activity phase preference. *Am. J. Physiol.*, 2000, 278: R1385–R1389.
- Kattler, H., Dijk, D. J. and Borbély, A. Effect of unilateral sensory stimulation prior to sleep on the sleep EEG in humans. *J. Sleep Res.*, 1994, 3: 159–164.
- Koorengevel, K. M., Beersma, D. G. M., Den Boer, J. A. and Van Den Hoofdakker, R. H. A forced desynchrony study of circadian pacemaker characteristics in seasonal affective disorder. *J. Biol. Rhythms*, 2002a, 17: 463–475.
- Koorengevel, K. M., Beersma, D. G. M., Den Boer, J. A. and Van Den Hoofdakker, R. H. Sleep in seasonal affective disorder patients in forced desynchrony: an explorative study. *J. Sleep Res.*, 2002b, 11: 347–356.
- Kräuchi, K. and Wirz-Justice, A. Circadian clues to sleep onset mechanisms. *Neuropsychopharmacology*, 2001, 25: S92–S96.
- Krueger, J. M. and Obal, F. A neuronal group-theory of sleep function. *J. Sleep Res.*, 1993, 2: 63–69.
- Krueger, J. M. and Tononi, G. Local use-dependent sleep; synthesis of the new paradigm. *Curr. Top. Med. Chem.*, 2011, 11: 2490–2492.
- Landolt, H. P. Sleep homeostasis: a role for adenosine in humans? *Biochem. Pharmacol.*, 2008, 75: 2070–2079.
- Landsness, E. C., Goldstein, M. R., Peterson, M. J., Tononi, G. and Benca, R. M. Antidepressant effects of selective slow wave sleep deprivation in major depression: a high-density EEG investigation. *J. Psychiatr. Res.*, 2011, 45: 1019–1026.
- Lazar, A. S., Lazar, Z. I. and Dijk, D. J. Circadian regulation of slow waves in human sleep: topographical aspects. *NeuroImage*, 2015, 116: 123–134.
- Legates, T. A., Altimus, C. M., Wang, H. *et al.* Aberrant light directly impairs mood and learning through melanopsin-expressing neurons. *Nature*, 2012, 491: 594–598.

- Leproult, R., Holmback, U. and Van Cauter, E. Circadian misalignment augments markers of insulin resistance and inflammation, independently of sleep loss. *Diabetes*, 2014, 63: 1860–1869.
- Levy, O., Dayan, T. and Kronfeld-Schor, N. The relationship between the Golden spiny mouse circadian system and its diurnal activity: an experimental field enclosures and laboratory study. *Chronobiol. Int.*, 2007, 24: 599–613.
- Martiny, K., Refsgaard, E., Lund, V. et al. A 9-week randomized trial comparing a chronotherapeutic intervention (Wake and Light Therapy) to exercise in major depressive disorder patients treated with duloxetine. *J. Clin. Psychiatry*, 2012, 73: 1234–1242.
- Meijer, J. H., Watanabe, K., Schaap, J., Albus, H. and Detari, L. Light responsiveness of the suprachiasmatic nucleus: long-term multi-unit and single-unit recordings in freely moving rats. *J. Neurosci.*, 1998, 18: 9078–9087.
- Mistlberger, R. E., Landry, G. J. and Marchant, E. G. Sleep deprivation can attenuate light-induced phase shifts of circadian rhythms in hamsters. *Neurosci. Lett.*, 1997, 238: 5–8.
- Mohawk, J. A., Baer, M. L. and Menaker, M. The methamphetamine-sensitive circadian oscillator does not employ canonical clock genes. *Proc. Natl Acad. Sci. USA*, 2009, 106: 3519–3524.
- Oliet, S. H. and Poulain, D. A. Adenosine-induced presynaptic inhibition of IPSCs and EPSCs in rat hypothalamic supraoptic nucleus neurons. *J. Physiol.*, 1999, 520: 815–825.
- O'Neill, J. S. and Reddy, A. B. Circadian clocks in human red blood cells. *Nature*, 2011, 469: 498–504.
- Pflug, B. and Töle, R. Therapy of endogenous depressions using sleep deprivation. Practical and theoretical consequences. *Nervenarzt*, 1971, 42: 117–124.
- Phillips, A. J. K., Czeisler, C. A. and Klerman, E. B. Revisiting spontaneous internal desynchrony using a quantitative model of sleep physiology. *J. Biol. Rhythms*, 2011, 26: 441–453.
- Plante, D. T., Landsness, E. C., Peterson, M. J. et al. Sex-related differences in sleep slow wave activity in major depressive disorder: a high-density EEG investigation. *BMC Psychiatry*, 2012, 12: 146.
- Porkka-Heiskanen, T., Strecker, R. E., Thakkar, M., Bjorkum, A. A., Greene, R. W. and McCarley, R. W. Adenosine: a mediator of the sleep-inducing effects of prolonged wakefulness. *Science*, 1997, 276: 1265–1268.
- Roecklein, K., Wong, P., Ernecoff, N. et al. The post illumination pupil response is reduced in seasonal affective disorder. *Psychiatry Res.*, 2013, 210: 150–158.
- Rosenthal, N. E., Sack, D. A., Gillin, J. C. et al. Seasonal affective disorder – a description of the syndrome and preliminary findings with light therapy. *Arch. Gen. Psychiatry*, 1984, 41: 72–80.
- Rusterholz, T. and Achermann, P. Topographical aspects in the dynamics of sleep homeostasis in young men: individual patterns. *BMC Neurosci.*, 2011, 1284: 1–10.
- Sahlem, G. L., Kalivas, B., Fox, J. B. et al. Adjunctive triple chronotherapy (combined total sleep deprivation, sleep phase advance, and bright light therapy) rapidly improves mood and suicidality in suicidal depressed inpatients: an open label pilot study. *J. Psychiatr. Res.*, 2014, 59: 101–107.
- Sato, T. and Kawamura, H. Circadian rhythms in multiple unit activity inside and outside the suprachiasmatic nucleus in the diurnal chipmunk *Eutamias sibiricus*. *Neurosci. Res.*, 1984, 1: 45–52.
- Schmidt, C., Collette, F., Leclercq, Y. et al. Homeostatic sleep pressure and responses to sustained attention in the suprachiasmatic area. *Science*, 2009, 324: 516–519.
- Spindelegger, C., Stein, P., Wadsak, W. et al. Light-dependent alteration of serotonin-1A receptor binding in cortical and subcortical limbic regions in the human brain. *World J. Biol. Psychiatry*, 2012, 13: 413–422.
- Stephan, F. K., Swann, J. M. and Sisk, C. L. Entrainment of circadian rhythms by feeding schedules in rats with suprachiasmatic lesions. *Behav. Neural Biol.*, 1979, 25: 545–554.
- Terman, M. and Terman, J. S. Controlled trial of naturalistic dawn simulation and negative air ionization for seasonal affective disorder. *Am. J. Psychiatry*, 2006, 163: 2126–2133.
- Thaiss, C. A., Zeevi, D., Levy, M. et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell*, 2014, 159: 514–529.
- Tobler, I. Effect of forced locomotion on the rest–activity cycle of the cockroach. *Behav. Brain Res.*, 1983, 8: 351–360.
- Tobler, I. and Stalder, J. Rest in the scorpion – a sleep-like state. *J. Comp. Physiol. A.*, 1988, 163: 227–235.
- Tobler, I., Borbély, A. A. and Groos, G. The effect of sleep deprivation on sleep in rats with suprachiasmatic lesions. *Neurosci. Lett.*, 1983, 42: 49–54.
- Tomotani, B. M., Flores, D. E. F. L., Tachinardi, P., Paliza, J. D., Oda, G. A. and Valentinuzzi, V. S. Field and laboratory studies provide insights into the meaning of day-time activity in a subterranean rodent (*Ctenomys aff. knighti*), the Tuco-Tuco. *PLoS One*, 2012, 7: 1–8.
- Tononi, G. and Cirelli, C. Sleep function and synaptic homeostasis. *Sleep Med. Rev.*, 2006, 10: 49–62.
- Tononi, G. and Cirelli, C. Sleep and the price of plasticity: from synaptic and cellular homeostasis to memory consolidation and integration. *Neuron*, 2014, 81: 12–34.
- Trachsel, L., Edgar, D. M., Seidel, W. F., Heller, H. C. and Dement, W. C. Sleep homeostasis in suprachiasmatic nuclei-lesioned rats – effects of sleep deprivation and triazolam administration. *Brain Res.*, 1992, 589: 253–261.
- Van der Vinne, V., Riede, S., Gorter, J. A. et al. Cold and hunger induce diurnality in a nocturnal mammal. *Proc. Natl Acad. Sci. USA*, 2014, 111: 15256–15260.
- Van Diepen, H. C., Ramkisoensing, A., Peirson, S. N., Foster, R. G. and Meijer, J. H. Irradiance encoding in the suprachiasmatic nuclei by rod and cone photoreceptors. *FASEB J.*, 2013, 27: 4204–4212.
- Van Diepen, H. C., Lucassen, E. A., Yassenkov, R. et al. Caffeine increases light responsiveness of the mouse circadian pacemaker. *Eur. J. Neurosci.*, 2014, 40: 3504–3511.
- Vandewalle, G., Hebert, M., Beaulieu, C. et al. Abnormal hypothalamic response to light in seasonal affective disorder. *Biol. Psychiatry*, 2011, 70: 954–961.
- Vansteensel, M. J., Michel, S. and Meijer, J. H. Organization of cell and tissue circadian pacemakers: a comparison among species. *Brain Res. Rev.*, 2008, 58: 18–47.
- Vyazovskiy, V. V., Achermann, P. and Tobler, I. Sleep homeostasis in the rat in the light and dark period. *Brain Res. Bull.*, 2007, 74: 37–44.
- Wehr, T. A. and Wirz-Justice, A. Internal coincidence model for sleep deprivation and depression. In: W. P. Koella (Ed) *Sleep 1980*. S. Karger, Basel, 1981: 26–33.
- Werth, E., Achermann, P. and Borbély, A. A. Brain topography of the human sleep EEG: antero-posterior shifts of spectral power. *NeuroReport*, 1996a, 8: 123–127.
- Werth, E., Dijk, D. J., Achermann, P. and Borbély, A. A. Dynamics of the sleep EEG after an early evening nap: experimental data and simulations. *Am. J. Physiol.*, 1996b, 271: R501–R510.
- Wirz-Justice, A. and Van Den Hoofdakker, R. H. Sleep deprivation in depression: what do we know, where do we go? *Biol. Psychiatry*, 1999, 46: 445–453.
- Wirz-Justice, A., Kräuchi, K., Brunner, D. P. et al. Circadian rhythms and sleep regulation in seasonal affective disorder. *Acta Neuropsychiatr.*, 1995, 7: 41–43.
- Wirz-Justice, A., Cajochen, C. and Nussbaum, P. A schizophrenic patient with an arrhythmic circadian rest–activity cycle. *Psychiatry Res.*, 1997, 73: 83–90.
- Wirz-Justice, A., Benedetti, F. and Terman, M. *Chronotherapeutics for Affective Disorders. A Clinician's Manual for Light and Wake Therapy*, 2nd edn. S. Karger AG, Basel, 2013.

- Wu, J. C., Kelsoe, J. R., Schachat, C. *et al.* Rapid and sustained antidepressant response with sleep deprivation and chronotherapy in bipolar disorder. *Biol. Psychiatry*, 2009, 66: 298–301.
- Wyatt, J. K., Ritz-De Cecco, A., Czeisler, C. A. and Dijk, D. J. Circadian temperature and melatonin rhythms, sleep, and neurobehavioral function in humans living on a 20-h day. *Am. J. Physiol.*, 1999, 277: R1152–R1163.
- Zarrinpar, A., Chaix, A., Yooseph, S. and Panda, S. Diet and feeding pattern affect the diurnal dynamics of the gut microbiome. *Cell Metab.*, 2014, 20: 1006–1017.
- Zavada, A., Strijkstra, A. M., Boerema, A. S., Daan, S. and Beersma, D. G. M. Evidence for differential human slow-wave activity regulation across the brain. *J. Sleep Res.*, 2009, 18: 3–10.